

DFTT 24/98
 UWThPh-1998-23
 hep-ph/9805387

Four-neutrino mixing and long-baseline neutrino oscillation experiments*

S.M. Bilenky

Joint Institute for Nuclear Research, Dubna, Russia

C. Giunti

*INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino,
 Via P. Giuria 1, I-10125 Torino, Italy*

W. Grimus

*Institute for Theoretical Physics, University of Vienna,
 Boltzmannngasse 5, A-1090 Vienna, Austria*

May 1998

Abstract

We consider the two schemes with four massive neutrinos which are compatible with the results of all neutrino oscillation experiments. We show that in these two schemes the probabilities of $\bar{\nu}_e$ disappearance and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance in long-baseline experiments are strongly suppressed.

The investigation of neutrino properties, in particular of neutrino masses and mixing, is one of the most important problems in today's high-energy physics and many experiments are dedicated to it. Among the numerous existing experimental results there are three positive indications which come from neutrino oscillation experiments. Neutrino oscillations can occur only if neutrinos are massive particles, if their masses are different and if neutrino mixing is realized in nature [1]. In this case, the left-handed flavor neutrino fields $\nu_{\alpha L}$ ($\alpha = e, \mu, \tau$) are superpositions of the left-handed components ν_{kL} ($k = 1, \dots, n$) of the fields of neutrinos with definite masses m_k :

$$\nu_{\alpha L} = \sum_{k=1}^n U_{\alpha k} \nu_{kL}, \quad (1)$$

*Talk presented by C. Giunti at the XXXIIInd *Rencontres de Moriond: Electroweak Interactions and Unified Theories*, Les Arcs 1800 (France), March 14-21 1998.

where U is a unitary mixing matrix. The general expression for the probability of $\nu_\alpha \rightarrow \nu_\beta$ transitions in vacuum is

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \left| \sum_{k=1}^n U_{\beta k} \exp \left(-i \frac{\Delta m_{k1}^2 L}{2E} \right) U_{\alpha k}^* \right|^2, \quad (2)$$

where Δm_{k1}^2 is the mass-squared difference $m_k^2 - m_1^2$, L is the distance between the neutrino source and detector and E is the neutrino energy.

The three experimental indications in favor of neutrino oscillations come from the solar neutrino problem [2], the atmospheric neutrino anomaly [3] and the results of the LSND experiment [4]. The solar neutrino deficit can be explained by transitions of ν_e 's into other states due to a mass-squared difference $\Delta m_{\text{sun}}^2 \sim 10^{-5} \text{ eV}^2$ in the case of resonant MSW transitions or $\Delta m_{\text{sun}}^2 \sim 10^{-10} \text{ eV}^2$ in the case of vacuum oscillations. The atmospheric neutrino anomaly can be explained by transitions of ν_μ 's into other states due to a mass-squared difference $\Delta m_{\text{atm}}^2 \sim 5 \times 10^{-3} \text{ eV}^2$. The recent results of the first reactor long-baseline (LBL) neutrino oscillation experiment CHOOZ [5] exclude an explanation of the atmospheric neutrino anomaly through $\nu_\mu \rightleftharpoons \nu_e$ oscillations, leaving the two possibilities: $\nu_\mu \rightarrow \nu_\tau$ or $\nu_\mu \rightarrow \nu_s$, where ν_s is a sterile neutrino. Finally, the LSND experiment found indications in favor of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations with a mass-squared difference in the range $0.3 \text{ eV}^2 \lesssim \Delta m_{\text{LSND}}^2 \lesssim 2.2 \text{ eV}^2$, which takes into account the negative results of other short-baseline (SBL) experiments [6]. Therefore, the three experimental indications in favor of neutrino oscillations imply that there are three independent mass-squared differences and the number of massive neutrinos is bigger than three. In the following we consider the simplest possibility of existence of four massive neutrinos. In this case, besides the three light flavor neutrinos ν_e, ν_μ, ν_τ that contribute to the invisible width of the Z -boson measured with high accuracy by LEP experiments, there is a light sterile flavor neutrino ν_s that is a $\text{SU}(2)_L$ singlet and does not take part in the standard weak interactions.

Almost two years ago we have shown [7] that among all the possible four-neutrino mass spectra only two [8] are compatible with the results of all neutrino oscillation experiments:

$$(A) \quad \underbrace{m_1 < m_2}_{\text{atm}} \ll \underbrace{m_3 < m_4}_{\text{sun}}, \quad (B) \quad \underbrace{m_1 < m_2}_{\text{sun}} \ll \underbrace{m_3 < m_4}_{\text{atm}}. \quad (3)$$

LSND LSND

In these two schemes the four neutrino masses are divided in two pairs of close masses separated by a gap of about 1 eV, which provides the mass-squared difference $\Delta m^2 \equiv \Delta m_{41}^2 \equiv m_4^2 - m_1^2$ that is relevant for the oscillations observed in the LSND experiment. In scheme A, $\Delta m_{21}^2 \equiv m_2^2 - m_1^2$ is relevant for the explanation of the atmospheric neutrino anomaly and $\Delta m_{43}^2 \equiv m_4^2 - m_3^2$ is relevant for the suppression of solar ν_e 's. In scheme B, the roles of Δm_{21}^2 and Δm_{43}^2 are reversed.

Let us define the quantities c_α , with $\alpha = e, \mu$, in the schemes A and B as

$$(A) \quad c_\alpha \equiv \sum_{k=1,2} |U_{\alpha k}|^2, \quad (B) \quad c_\alpha \equiv \sum_{k=3,4} |U_{\alpha k}|^2. \quad (4)$$

The results of all neutrino oscillation experiments are compatible with the schemes A and B if [7]

$$c_e \leq a_e^0 \quad \text{and} \quad c_\mu \geq 1 - a_\mu^0, \quad (5)$$

where

$$a_\alpha^0 \equiv \frac{1}{2} \left(1 - \sqrt{1 - B_{\alpha;\alpha}^0} \right) \quad (\alpha = e, \mu) \quad (6)$$

and $B_{\alpha;\alpha}^0$ is the upper bound for the amplitude of $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha$ oscillations obtained from the exclusion plots of SBL reactor and accelerator disappearance experiments. Therefore, the quantities a_e^0 and a_μ^0 depend on Δm^2 . The exclusion curves obtained in the Bugey reactor experiment and in the CDHS and CCFR accelerator experiments [9] imply that both a_e^0 and a_μ^0 are small quantities [10]: $a_e^0 \lesssim 4 \times 10^{-2}$ and $a_\mu^0 \lesssim 2 \times 10^{-1}$ for any value of Δm^2 in the range $0.3 \lesssim \Delta m^2 \lesssim 10^3 \text{ eV}^2$.

The smallness of c_e in both schemes A and B is a consequence of the solar neutrino problem [7]. It implies that the electron neutrino has a small mixing with the neutrinos whose mass-squared difference is responsible for the oscillations of atmospheric neutrinos (*i.e.*, ν_1, ν_2 in scheme A and ν_3, ν_4 in scheme B). Hence, the transition probability of electron neutrinos and antineutrinos into other states in atmospheric and long-baseline experiments is suppressed. Indeed, it can be shown [11] that the transition probabilities of electron neutrinos and antineutrinos into all other states are bounded by

$$1 - P_{\bar{\nu}_e \rightarrow \bar{\nu}_e}^{(\text{LBL})} \leq a_e^0 (2 - a_e^0) . \quad (7)$$

The curve corresponding to this limit obtained from the 90% CL exclusion plot of the Bugey experiment is shown in Fig.1 (solid line). The shadowed region in Fig.1 corresponds to the range of Δm^2 allowed at 90% CL by the results of the LSND and all the other SBL experiments. The upper bound for the transition probability of $\bar{\nu}_e$'s into other states obtained from the recently published exclusion plot of the CHOOZ [5] experiment is shown in Fig.1 (dash-dotted line). One can see that the CHOOZ upper bound agrees with that obtained from Eq.(7). In Fig.1 we have also drawn the curve corresponding to the expected final sensitivity of the CHOOZ experiment (dash-dot-dotted line). One can see that the LSND-allowed region imply an upper bound for $1 - P_{\bar{\nu}_e \rightarrow \bar{\nu}_e}^{(\text{LBL})}$ of about 5×10^{-2} , which is smaller than the expected final sensitivity of the CHOOZ experiment.

The probability of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ transitions in LBL experiments is bounded by [11]

$$P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e}^{(\text{LBL})} \leq \min \left(a_e^0 (2 - a_e^0) , a_e^0 + \frac{1}{4} A_{\mu;e}^0 \right) , \quad (8)$$

where $A_{\mu;e}^0$ is the upper bound for the amplitude of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ transitions measured in SBL experiments with accelerator neutrinos. The bound obtained with Eq.(8) from the 90% CL exclusion plots of the Bugey experiment and of the BNL E734, BNL E776 and CCFR experiments [12] is depicted by the dashed line in Fig.2. The solid line in Fig.2 shows the upper bound on $P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e}^{(\text{LBL})}$ taking into account matter effects. The expected sensitivities of the K2K long-baseline accelerator neutrino experiment [13] is indicated in Fig.2 by the dash-dotted vertical line. The shadowed region in Fig.2 corresponds to the range of Δm^2 allowed at 90% CL by the results of the LSND and all the other SBL experiments. It can be seen that the results of SBL experiments indicate an upper bound for $P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e}^{(\text{LBL})}$ smaller than 4×10^{-2} and smaller than the expected sensitivity of the K2K experiment.

In conclusion, we have shown that in the four-neutrino schemes A and B which are compatible with the results of all neutrino oscillation experiments, the probabilities of $\bar{\nu}_e$ disappearance and




Figure 1




Figure 2

$\nu_{\mu}^{(-)} \rightarrow \nu_e^{(-)}$ appearance in LBL experiments are severely constrained (on the other hand, the channels $\nu_{\mu}^{(-)} \rightarrow \nu_{\mu}^{(-)}$ and $\nu_{\mu}^{(-)} \rightarrow \nu_{\tau}^{(-)}$ are not constrained at all). The two schemes A and B have identical implications for neutrino oscillation experiments, but very different implications for neutrinoless double- β decay experiments and for tritium β -decay experiments. Indeed, in scheme A

$$|\langle m \rangle| \leq m_4 \quad \text{and} \quad m(^3\text{H}) \simeq m_4, \quad (9)$$

whereas in scheme B

$$|\langle m \rangle| \leq a_e^0 m_4 \ll m_4 \quad \text{and} \quad m(^3\text{H}) \ll m_4, \quad (10)$$

where $\langle m \rangle = \sum_{i=1}^4 U_{ei}^2 m_i$ is the effective Majorana mass that determines the matrix element of neutrinoless double- β decay and $m(^3\text{H})$ is the neutrino mass measured in tritium β -decay experiments. Therefore, in scheme B $|\langle m \rangle|$ and $m(^3\text{H})$ are smaller than the expected sensitivity of the next generation of neutrinoless double- β decay and tritium β -decay experiments. The observation of a positive signal in these experiments would be an indication in favor of scheme A.

References

References

- [1] See S.M. Bilenky and B. Pontecorvo, Phys. Rep. **41**, 225 (1978).
- [2] B.T. Cleveland *et al.*, Nucl. Phys. B (P.S.) **38**, 47 (1995); K.S. Hirata *et al.*, Phys. Rev. D **44**, 2241 (1991); GALLEX Coll., Phys. Lett. B **388**, 384 (1996); SAGE Coll., Phys. Rev. Lett. **77**, 4708 (1996); M. Nakahata, Talk presented at the APCTP Workshop: *Pacific Particle Physics Phenomenology*, Seoul, Korea, October 31 – November 2, 1997 (http://www-sk.icrr.u-tokyo.ac.jp/doc/sk/pub/pub_sk.html).
- [3] Y. Fukuda *et al.*, Phys. Lett. B **335**, 237 (1994); R. Becker-Szendy *et al.*, Nucl. Phys. B (P.S.) **38**, 331 (1995); W.W.M. Allison *et al.*, Phys. Lett. B **391**, 491 (1997); Y. Fukuda *et al.*, hep-ex/9803006.
- [4] C. Athanassopoulos *et al.*, Phys. Rev. Lett. **77**, 3082 (1996).
- [5] M. Apollonio *et al.*, Phys. Lett. B **420**, 397 (1998).
- [6] For a review see J. Brunner, Fortsch. Phys. **45**, 343 (1997).
- [7] S.M. Bilenky, C. Giunti and W. Grimus, Proc. of *Neutrino96*, Helsinki, June 1996, edited by K. Enqvist *et al.*, p.174 (World Scientific, 1997); Eur. Phys. J. C **1**, 247 (1998).
- [8] See also N. Okada and O. Yasuda, Int. J. Mod. Phys. A **12**, 3669 (1997).
- [9] B. Achkar *et al.*, Nucl. Phys. B **434**, 503 (1995); F. Dydak *et al.*, Phys. Lett. B **134**, 281 (1984); I.E. Stockdale *et al.*, Phys. Rev. Lett. **52**, 1384 (1984).
- [10] See Fig.1 of S.M. Bilenky, A. Bottino, C. Giunti and C.W. Kim, Phys. Rev. D **54**, 1881 (1996).
- [11] S.M. Bilenky, C. Giunti and W. Grimus, Phys. Rev. D **57** (1998) 1920.
- [12] L.A. Ahrens *et al.*, Phys. Rev. D **36**, 702 (1987); L. Borodovsky *et al.*, Phys. Rev. Lett. **68**, 274 (1992); A. Romosan *et al.*, Phys. Rev. Lett. **78**, 2912 (1997).
- [13] Y. Oyama (for the K2K collaboration), hep-ex/9803014.